Helianthemum sicanorum (Cistaceae), a new species from Sicily

by

Salvatore Brullo, Gian Pietro Giusso del Galdo & Saverio Sciandrello

Dipartimento di Botanica, via A. Longo 19, I - 95125 Catania, Italy
brullo@dipbot.unict.it

Abstract

Brullo, S., Giusso del Galdo, G.P. & Sciandrello, S. 2007. *Helianthemum sicanorum* (Cistaceae), a new species from Sicily. *Anales Jard. Bot. Madrid* 64(1): 47-53.

Helianthemum sicanorum, an endemic species occurring on calcareous marls near Gela (S Sicily), is described and illustrated. Due to its habit and flower morphology, *H. sicanorum* shows close relationships with *H. kahiricum* Del., circumscribed to the semi-arid habitats of northern Africa and the Middle East. *H. sicanorum* is quite rare and localized on steep slopes facing the sea, where it is a member of thermophilous garigues of the Cisto-Micromerietea class.

Keywords: Cistaceae, endemism, *Helianthemum*, phytosociology, Sicily, taxonomy.

Resumen

Brullo, S., Giusso del Galdo, G.P. & Sciandrello, S. 2007. *Helianthemum sicanorum* (Cistaceae), una nueva especie de Sicilia. *Anales Jard. Bot. Madrid* 64(1): 47-53 (en inglés).

Se describe e ilustra la nueva especie *Helianthemum sicanorum*, que crece sobre margas calizas cerca de Gela, en el sur de Sicilia. Por su hábito y morfología floral, *H. sicanorum* está relacionado con *H. kahiricum* Del., especie propia de ambientes subdesérticos del norte de África y Asia Menor. *H. sicanorum* es muy raro, y está limitado a laderas abruptas que miran al mar, donde crece en garrigas termófilas de la clase Cisto-Micromerietea.

Palabras clave: Cistaceae, endemismo, fitosociología, *Helianthemum*, Sicilia, taxonomía.

Introduction

The coastal belt of southern Sicily represents a very important area from the phytogeographical point of view. In fact, this part of the island is characterized by the occurrence of several endemic or rare species, mostly circumscribed to this territory (Giusso del Galdo & Sciandrello, 2003). During floristic investigations along the rocky coast of "Torre Manfredi" (Gela), a small population of a really peculiar and hitherto uncollected *Helianthemum* was found (Proctor & Heywood, 1968; Pignatti, 1982). Based on investigations carried out on herbarium material (CAT, FI, RO) and literature data, this plant is closely related to *H. kahiricum* Delile, a xerophyte distributed in arid and semi-arid habitats of N Africa and the Middle East, but not recorded for any European country. However, the Sicilian population, differing from the typical of *H. kahiricum* in many morphological features chiefly concerning the leaves, inflorescence,

flowers and fruits, is here described as a species new to science.

Materials and methods

The morphological study of *H. sicanorum* was based on living plants coming from the type locality, while *H. kahiricum* was examined using herbarium specimens (CAT, FI) and the measurements of the flowers were taken on dry hydrated specimens.

Helianthemum sicanorum Brullo, Giusso & Sciandrello, **sp. nov.** (Figs. 1, 2)

Ab Helianthemum kahiricum differt caulibus adpresse lanuginosis pilisque simplicibus 0,1-0,3(0,4) mm longis, foliis etiam ita indutis atque linearibus aut lineari-oblongeolatis, petiolis 0,3-0,8 mm longis, inflorescentia maxime 10-flora, bracteis 1,5-2 mm longis, pedicellis 5-7 mm longis, calyce albo-sericeo pilis simplicibus adpres-

Anales del Jardín Botánico de Madrid 64(1): 47-53, enero-junio 2007. ISSN: 0211-1322

sisque atque 0,1-0,2 mm longis, sepalis exterioribus subulatis apiceque rotundatis atque 0,1-0,2 mm longis, sepalis interioribus 3-nervibus apiceque obtusiusculis atque 4-5,5 mm longis, corolla calycem excedenti vel aequanti, petalis 4-5 × 2-2,5 mm longis latisque, filamentis staminalibus 1-1,6 mm longis, anthera 0,4-0,5 mm longa, ovario 1-1,2 mm longo, stylo 1,4-1,6 mm longo, stigmate papillis multis brevibusque (0,01 mm longis) cooperto, capsula 3,5 × 1,8-2 mm longa lataque atque pilis 0,1-0,3 mm longis tecta seminibusque brunneo-rubrescentibus atque 1-1,1 mm longis.

Small shrub 20-30 cm tall, stems laxly caespitose, branched from the base, erect to ascending, with stellate hairs and shorter simplex hairs 0.1-0.3(0.4) mm long. Indumentum tomentose-canescens and appressed in the stem and leaves. Leaves linear to linear-oblongate 3.5-10 mm long, 1-2(3) mm wide, strongly revolute at the margins with petiole 0.3-0.8 mm long. Stipules ovate-lanceolate, longer than the petiole, 1-1.5 mm long, 0.4-0.6 mm wide. Inflorescence one-sided, 5-10 flowered. Bracts lanceolate, 1.5-2 mm long. Pedicels exceeding or equalling the calyx, 5-7 mm long, patent or deflexed. Calyx white-sericeous, green-purplish, with simplex hairs 0.1-0.2 mm long. Outer sepals subulate 2-3 mm long, rounded at the apex; inner sepals oblong-ovate, lightly obtuse at apex, 4-5.5 mm long, 2.8-3.2 mm wide, 3-nerved. Corolla exceeding or equalling the calyx, with petals yellow 4-5 mm long, 2-2.5 mm wide, many nerved, often not opening (cleistogamous flowers). Stamens numerous with filament 1-1.6 mm long and anther 0.4-0.5 mm long. Ovary ovoid, densely hairy, 1-2 mm long. Style white, curved 1.4-1.6 mm long. Stigma sulcate, sub-plane, covered by numerous short papillae (ca. 0.01 mm). Capsule ellipsoid, 3.5 mm long, 1.8-2 mm wide, densely pilose, with hairs 0.1-0.3 mm long. Seeds brown-reddish, minutely punctate, 1-1.1 mm long.

Holotype: Sicily, Gela, litorale roccioso presso Torre Manfria, 30-VII-2005, *Sciandrello s.n.* (CAT; isotypes, CAT, FI, MA).

Etymology. The specific epithet refers to “Sicani”, an ancient civilization settled in Sicily from the second millennium to the fifth century B.C.

Distribution. This species is known only for a small population of no more than 50 individuals, circumscribed to Torre Manfria near Gela, (S Sicily), along a narrow strip of rocky coast.

Habitat. *Helianthemum sicanorum* grows on Pliocene marly limestones facing the sea, where it colonizes steep and hardly reachable slopes (Fig. 3). From the bioclimatic viewpoint the area at issue belongs to the bioclimatic belt with a lower thermo-

Fig. 2. *Helianthemum sicanorum*: inflorescence (from the type locality, July 30th 2005).

mediterranean thermotype and upper semiarid ombrottype (Brullo & al. 1996). This species plays an important physiognomic role within the thermophilous garigues characterized by undersized chamaephytes linked to really arid habitats. In particular, it is a member of the *Diplofaxio crassifoliae*-*Reaumurieta* *vermiculatae* (Table 1), a xeric plant community described by Brullo & al. (2000) belonging to the Cisto-Micromerietea class. Within this association, some North African species (*Reaumuria vermiculata* L., *Diplofaxis crassifolia* (Rafin.) DC., etc.), usually rare for the Sicilian flora, are frequent too.

Phenology: flowering February to July; fruiting April to August.

Conservation status. At present, *H. sicanorum* is a threatened species at high risk of extinction as a result of the low number of individuals (less than 50) circumscribed to a narrow stand close to the sea. Therefore, since this species is seriously endangered by several human activities (such as tourist facilities, fire, urban sprawl, agricultural exploitation, and so on), we deem it should be added to the Regional Red List of Threatened Species as critically endangered (CR).

Fig. 3. Habitat of *Helianthemum sicanorum*. Locality of holotype, within garigue of *Diplofaxia crassifoliae*-*Reaumurieta vermiculatae* (Cisto-Micromerietea), July 30th 2005.

Based on the criteria adopted by IUCN (2001, 2003, 2005), it is proposed its inclusion in the following category: CR B1a, B2a, D.

Taxonomic remarks. According to the classification proposed by Grosser (1903) for the genus *Helianthemum*, this new species has to be referred to the Sect. *Eriocarpum* Dunal. This section groups deserticolous shrubs characterized by unbranched inflorescence, persistent and pungent when dried, stamens 15-20, style ascending to geniculate, capsule rounded-trigone, pubescent to hirsute, flowers often cleistogamous with petals contorted at the apex, enveloping the capsule as a mitre. In its morphology, *Helianthemum sicanorum* is quite similar to *H. kabiricum*, a species widespread in the South Mediterranean countries, described and illustrated by Delile (1813) from northern Egypt. Both these taxa are small shrubs with a woody subspinescent habit and branches covered by dense whitish tomentum, but several different features allow them to be easily distinguished. In particular, *H. kabiricum* has a patent and longer indumentum with hairs

from 0.5-1.5 mm long, and linear lanceolate leaves up to 20 × 7 mm. The petiole is 0.5-2 mm long, the inflorescence up to 12-flowered, bracts 0.2-0.5 mm long, pedicels 2-5 mm long, and the calyx villous with simplex patent hairs 0.5 mm long. The outer sepals are lanceolate-subulate, acute at apex, 3-6 mm long, inner sepals 5-7 mm long, acute at apex, 4-nerved, corolla shorter than calyx with petals 2.7-3 mm long and 1.5-1.8 mm wide, stamen filaments 0.6-1 mm long, anther 0.3-0.4 mm long, ovary 1.2-1.4 mm long, style 0.8-1 mm long, stigma covered by few longer papillae (0.03-0.06 mm long), capsule 4-5 mm long and 2.3-2.5 mm wide, covered by hairs 0.3-0.5 mm long, and the seeds are dark-brown 1.2-1.3 mm long (Fig. 4). From the point of view of chorology, and based on the following authors [Bonnet & Barratte (1896), Grosser (1903), Muschler (1912), Pampanini (1914), Post (1932), Quézel & Santa (1963), Zohary (1972), Jafri (1977), Pottier-Alapetite (1979), Collenette (1985), Mouterde (1970), Migahid (1988), Feinbrun-Dothan & Danin (1998), Raynaud (1999) and Boulos (2000)], *H. kabi-*

Table 1. *Diploaxis crassifoliae* - *Reaumurieta vermiculatae* Brullo, Guarino & Ronsisvalle 2000.

Number of inventory	1	2	3
Surface (m ²)	20	10	10
Cover (%)	60	70	60
Exposition	S	S	S
Gradient (°)	50	50	50
Characteristic of association			
<i>Helianthemum sicanorum</i> Brullo, Giusso & Sciandrello	2	2	1
<i>Reaumuria vermiculata</i> L.	1	+	1
<i>Diploaxis crassifolia</i> (Raf.) DC.	1	+	+
Characteristic of Cisto-Ericion & Cisto-Micromerietea			
<i>Thymelaea hirsuta</i> (L.) Endl.	1	1	2
<i>Coridothymus capitatus</i> (L.) Rchb.f.	3	3	3
<i>Phagnalon rupestre</i> (L.) DC. ssp. <i>annoticum</i> (Jord. ex Burnat) Pignatti	1	1	2
<i>Teucrium capitatum</i> L.	.	1	1
Other species			
<i>Reichardia picroides</i> (L.) Roth	1	1	1
<i>Daucus gingidium</i> L.	3	2	1
<i>Lygeum spartum</i> L.	1	2	1
<i>Trachynia distachya</i> (L.) Link	+	+	1
<i>Asparagus horridus</i> L.	1	+	1
<i>Sedum sediforme</i> (Jacq.) Pau	2	1	2
<i>Artemisia arborescens</i> L.	+	+	+
<i>Capparis spinosa</i> L.	1	.	+
<i>Pistacia lentiscus</i> L.	+	.	+
<i>Salsola oppositifolia</i> Desf.	.	+	+
<i>Plantago macrorrhiza</i> Poir.	2	.	.
<i>Limbarda crithmoides</i> (L.) Dumort.	+	.	.
<i>Suaeda vera</i> Forssk. ex J.F. Gmel.	1	.	.
<i>Teucrium fruticans</i> L.	.	1	.
<i>Dactylis hispanica</i> Roth	.	+	.
<i>Valantia muralis</i> L.	+	.	.
<i>Prasium majus</i> L.	.	.	1

ricum is a Saharo-Arabian element usually growing in desertic or the semi-arid places of N Africa, Sinai, Palestine, Syria, Iraq and Saudi Arabia (Fig. 5). Because of this, it seems quite possible that *H. sicanorum* should be as a geographical vicariant of *H. kahiricum*, replacing it in Sicily.

In conclusion, *H. sicanorum* represents an endemic with thermo-xerophilous requirements, showing a punctiform distribution area, which is likely to be the result of the long geographic isolation of the Sicilian population from the southern Mediterranean ones.

Representative specimens

Helianthemum sicanorum

ITALY: **Sicily**, Gela, litorale roccioso presso Torre Manfria, 24-VII-2005, *Sciandrello s.n.* (CAT); *ibid.*, 8-I-2006, *Sciandrello s.n.* (CAT); *ibid.*, 4-IV-2006, *Brullo, Giusso & Sciandrello s.n.* (CAT).

Helianthemum kahiricum

EGYPT: Gebel Mokattan, 2-III-1898, *Marchesetti s.n.* (FI); *ibid.*, pr. Cairo, II-1898, *Marchesetti s.n.* (FI). Heluan, 25-II-1898, *Marchesetti s.n.* (FI). Wadi Hof, sandy soil, 25-II-1961, *Khattab s.n.* (FI). Bassa Tebeide, nei valloni e burroni dell'altipiano del Mokattan, verso il Cairo nonché nelle montagne della Regione

Fig. 4. *Helianthemum kahircum* (Libya, CAT): **A**, habit; **B**, detail of inflorescence; **C**, leaves (upper surface); **D**, leaves (below surface); **E**, leaves (transversal section); **F**, flower bud; **G**, closed flower; **H**, open calyx; **I**, fructified flower; **J**, terminal part of the outer sepal; **K**, open flower; **L**, petals; **M**, stamens; **N**, pistil; **O**, stigma; **P**, capsule; **Q**, seeds.

Fig. 5. Distribution map of *Helianthemum sicanorum* (star) and *H. kahiricum* (literature data empty circle and herbarium specimens full circle).

Sinaica, IV, *Figari* (FI). Summit Jebel Mokattan about 5 miles SE of Cairo, 4-IV, *Scott Elliot*, 3617 (FI). Sull'altopiano calcareo della Valle Nilotica (Mokattan e Gebel Ghioceseli), IV, *Figari* (FI). Prope Cairo, s.d., *Samaritani s.n.* (FI). Nel deserto di Alessandria che segue il litorale del deserto di Ramba e di Abuk, V-1867, *Figari s.n.* (FI). Comune in luoghi rupestri della Regione del Sinai, ed è pure del deserto montano d'Egitto, s.d., *Figari s.n.* (FI). LIBYA: Cirenaica, Msus, 20-III-1933, *Pampanini* 5172 (FI). *ibid.*, Amseat a sud di Bardia 24-III-1963, *Pampanini* 5173 (FI). *ibid.*, fra El Aghelia e Agedabia, 22-V-1937, *Servizi Agrari Pirenaica* 233 (FI). *ibid.*, Uadi Faregh, Maatan Giafar a sud di El Aghelia, 15-III-1933, *Pampanini* 5171 (FI); *ibid.*, Saniet El Hamar a sud-est di Agedabia, 12-III-1933, *Pampanini* 5169 (FI); *ibid.*, Gebel el Bab, presso il Marabutto, 18-V-1981, *Brullo & Furnari s.n.* (CAT); *ibid.*, Ajedabia, 19-V-1981, *Brullo & Furnari s.n.* (CAT); *ibid.*, 80 Kil. De Derna (Soud de Tobruk), 24-IV-1938, *Maire & Weiller* 178 (FI); *ibid.*, Martuba, a sud-est di Barce, U. el Bgor, 8-IV-1933, *Pampanini* 5180 (FI); *ibid.*, Uadi Faregh, Maaten Giofer, 8-IV-1934, *Pampanini & Pichi-Sermolli* 5181 (FI). Sirtica or., a sud di Gas es Sahabi, 20-IV-1928, *Krueger s.n.* (FI); *ibid.*, fra l'Uadi Feregh e Maatan Risam, 18-IV-1928, *Krueger s.n.* (FI). Marmarica, Porto Bardia, 20-VII-1926, *Krueger s.n.* (FI); *ibid.*, Tobruk, 1918, *Cassinea* (FI). Tripolitania, Taruna, Uadi Ksea, Ras Argobinam, 8-IV-1913, *Pampanini* 2448 (FI); *ibid.*, Mesellata, Msid di Cesellata, 9-IV-1913, *Pampanini* 2580 (FI); *ibid.*, colline a N-NW di Cussabat, 12-IV-1913, *Pampanini* 2951 (FI); *ibid.*, Cussabat, Ras Gelà, 13-IV-1913, *Pampanini* 3134 (FI); *ibid.*, Uadi Gherrim, 15-IV-1913, *Pampanini* 3265 (FI); *ibid.*, Garian, Uadi Garian, 29-IV-1913, *Pampanini s.n.* (FI); *ibid.*, Ras Bu-Ganus, 27-IV-1913, *Pampanini* 3983 (FI). TUNISIA: Entre Bir et Haguët et Bir Teura, 14-IV-1887, *Letourneux s.n.* (FI). In collibus aridis deserti ad Sidi Bout-Baba prope Gabes, 7-III-1854, *Kralik s.n.* (FI). ALGERIA: In lapidosis arenosis montis Melia prope Figiug, 1000 m, 17-IV-1932, *Maire s.n.* (FI). Bistra, in glareosis planit. et in rupibus montium "Djebel Maovia, Djebel Snia", V-1896, *Chevallier* 16 (FI); *ibid.*, sur les coteaux pierreux, 20-IV-1853, *Balansa* (FI). Rochers de la Region Latoricum Tiaut, 24-IV-1896, *Chauber s.n.* (FI).

Acknowledgements

The authors thank the curators of the herbaria of CAT and FI for the consultation of the *exsiccata*, and M. Láinz for correcting the Latin diagnosis. Financial support from the ECONET-COHASt project (INTERREG IIIB – ARCHIMED) is also gratefully acknowledged.

References

- Bonnet, E. & Barratte, G. 1896. *Catalogue raisonné des plantes vasculaires de la Tunisie*. Imprimerie Nationale. Paris.
- Boulos, L. 2000. *Flora of Egypt, Geraniaceae – Boraginaceae*, Vol. 2. Al Hadara Publishing. Cairo.
- Brullo, S., Guarino, R. & Ronsisvalle, G. 2000. La vegetazione del litorale di Manfria, presso Gela (Sicilia), area soggetta a vincolo archeologico. *Arch. Geobot.* 4(1): 91-107.
- Brullo, S., Scelsi, F., Siracusa, G. & Spampinato, G. 1996. Caratteristiche bioclimatiche della Sicilia. *Giorn. Bot. Ital.* 130(1): 177-185.
- Collenette, S. 1985. *An illustrated guide to the flowers of Saudi Arabia*. Scorpion Publishing Ltd. London.
- Delile, A.R. 1813. *Flore d'Égypte*. In: *Description de l'Égypte... Histoire naturelle. Tom. second*. Imprimerie Impériale. Paris.
- Feinbrun-Dothan, N. & Danin, A. 1998. *Analytical flora of Eretz-Israel*. 2nd ed. Cana Publishing House Ltd. Jerusalem.
- Giusso del Galdo, G. & Sciandrello, S. 2003. Contributo alla flora dei dintorni di Gela (Sicilia meridionale). *Abstracts 98° Congr. Soc. Bot. Ital.*: 235. Catania.
- Grosser, W. 1903. Cistaceae. In: H.G.A. Engler (ed.), *Pflanzenreich* 14 (IV. 193): 1-161. Breitkopf & Härtel. Leipzig.
- IUCN. 2001. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN Species Survival Commission. IUCN. Gland & Cambridge.
- IUCN. 2003. *Guidelines for Application of IUCN Red List Criteria at Regional Levels: Version 3.0*. IUCN Species Survival Commission. IUCN. Gland and Cambridge.
- IUCN. 2005. *Guidelines for Using the IUCN Red List, Categories and Criteria*. IUCN Standards and Petitions Subcommittee. Gland & Cambridge.
- Jafri, S.M.H. 1977. Cistaceae. In: Jafri, S.M.H. & El-Gadi, A. (eds.), *Flora of Libya* 48: 1-43. Al Faateh University. Tripoli.
- Migahid, A. M. 1988. *Flora of Saudi Arabia*. Third edition. Vol. 1. King Saud University Libraries. Riyadh.
- Mouterde, P. 1970. *Nouvelle flore du Liban et de la Syrie*. Vol. 2. Dar El-Machreq Éditeurs. Beyrouth.
- Muschler, R. 1912. *A manual flora of Egypt*. Vol.1. Friedlaender & Sohn. Berlin.
- Pampanini, R. 1914. *Plantae Tripolitanae ab auctore anno 1913 lectae et repertorium florum vascularis Tripolitaniae*. Stabilimento Pellas Luigi Caiti successore. Firenze.
- Pignatti, S. 1982. *Helianthemum* Miller. In: Pignatti, S. (ed.), *Flora d'Italia* 2: 125-130. Edagricole. Bologna.
- Post, G.E. 1932. *Flora of Syria, Palestine and Sinai*. Vol. 1. American Press. Beirut.
- Pottier-Alapetite, G. 1979. *Flore de la Tunisie, Angiospermes-Dicotyledones, Apetales-Dialypetales*, Vol. 1. Imprimerie officielle de la République Tunisienne. Tunis.
- Proctor, M.C.F. & Heywood, V.H., 1968. *Helianthemum* Miller. In: Tutin, T.G. & al. (eds.), *Flora Europea* 2: 286-291. University Press. Cambridge.
- Quézel, P. & Santa, S. 1963. *Nouvelle Flore de l'Algérie et des régions désertiques méridionales*, Vol. 2. Editions du Centre National de la Recherche Scientifique. Paris.
- Raynaud, C. 1999. Cistaceae. In: M. Fennane et al. (eds.), *Flore Pratique du Maroc. Manuel de détermination des plantes vasculaires*, 1: 302-326. Trav. Inst. Scient., Université Mohammed V, s. Bot. 36. Rabat.
- Zohary, M. 1972. *Flora Palaestina, Platanaceae to Umbelliferae*, Vol. 2. The Israel Academy of Sciences and Humanities. Jerusalem.

Associate Editor: G. López

Received: 9-X-2006

Accepted: 29-III-2007